

ON THE LACK OF STRONG O-LINE EXCESS IN THE COMA CLUSTER OUTSKIRTS FROM *SUZAKU*

YOH TAKEI¹, ERIC D. MILLER², JOEL N. BREGMAN³, SHUNSUKE KIMURA⁴, TAKAYA OHASHI⁵, KAZUHIISA MITSUDA⁴,
TAKAYUKI TAMURA⁴, NORIKO Y. YAMASAKI⁴, RYUICHI FUJIMOTO⁶

Accepted 2008 March 19

ABSTRACT

About half of the baryons in the local Universe are thought to reside in the so-called warm-hot intergalactic medium (WHIM) at temperatures of 0.1-10 million K. Thermal soft excess emission in the spectrum of some cluster outskirts that contains OVII and/or OVIII emission lines is regarded as evidence of the WHIM, although the origin of the lines is controversial due to strong Galactic and solar system foreground emission. We observed the Coma-11 field, where the most prominent thermal soft excess has ever been reported, with *Suzaku* XIS in order to make clear the origin of the excess. We did not confirm OVII or OVIII excess emission. The OVII and OVIII intensity in Coma-11 is more than 5σ below that reported before and we obtained 2σ upper limits of 2.8 and 2.9 photons $\text{cm}^{-2} \text{s}^{-1} \text{sr}^{-1}$ for O VII and O VIII, respectively. The intensities are consistent with those in another field (Coma-7) that we measured, and with other measurements in the Coma outskirts (Coma-7 and X Com fields with XMM-Newton). We did not confirm the spatial variation within Coma outskirts. The strong oxygen emission lines previously reported are likely due to solar wind charge exchange.

Subject headings: clusters: individual: Coma — intergalactic medium — large-scale structure of universe

1. INTRODUCTION

Based on N-body simulations of cosmological large-scale structure formation (e.g., Cen & Ostriker 2006), a significant (30–50%) fraction of baryons are thought to reside in a ‘warm-hot’ gaseous phase ($T = 10^{5-7}$ K), which is hard to detect with the instruments currently in operation. This warm-hot gas, whose density is 10^{-6} – 10^{-4} cm^{-3} , is called the warm-hot intergalactic medium (WHIM). Firm detection of the WHIM is important because it is the most promising candidate for the “missing baryons”; i.e., it is thought to explain the discrepancy between the baryon density observed in the local universe (Fukugita et al. 1998) and that in the distant universe (Rauch et al. 1997) or that calculated from the observed fluctuations of the cosmic microwave background (Spergel et al. 2007). Although $\sim 10\%$ of the baryons has been resolved as the WHIM in the temperature range $T = 10^{5-6}$ K through O VI absorption features observed with *FUSE* and *HST* (Danforth & Shull 2007), more baryons are thought to reside in a hotter phase at $T = 10^{6-7}$ K. The WHIM of this temperature may be detected via emission or absorption lines from highly ionized elements in X-ray spectra, such as O VII and O VIII.

Despite many observations so far, the existence of the WHIM in the hotter phase is not yet confirmed (see re-

view of Bregman 2007, and references therein). Studies of the WHIM using absorption lines only lead to one disputed detection toward Mkn 421 (Nicastrro et al. 2005), a result which was not confirmed later and was thus questioned (Rasmussen et al. 2007; Kaastra et al. 2006). Possible detections of emission from the WHIM have been reported in cluster outskirts, as a soft X-ray ($E \lesssim 0.5 \text{ keV}$) excess containing O VII and O VIII lines (Kaastra et al. 2003; Finoguenov et al. 2003). The origin of O line emission is, however, controversial due to strong Galactic and solar system foreground emission, which has a spectral shape similar to the WHIM (Bonamente et al. 2005; Bregman & Lloyd-Davies 2006). In particular, X-rays induced by solar wind charge exchange (SWCX) could produce serious contamination because of its high variability (e.g., Cravens 2000; Fujimoto et al. 2007). Note that there has been no detection of redshifted emission lines from the WHIM so far.

The most prominent thermal soft excess ever reported is in the Coma outskirts, especially in the Coma-11 field (Finoguenov et al. 2003, hereafter FBH03). FBH03 also observed spatial variation within the Coma outskirts; Coma-7 shows roughly five times smaller intensity than Coma-11. Not only O lines, but also an EUV excess (Bonamente et al. 2003) and marginal detection of Ne IX emission and absorption lines (Takei et al. 2007, hereafter T07) are reported in the Coma outskirts. These results may support the existence of the WHIM.

We observed the Coma-11 and Coma-7 fields with the XIS (Koyama et al. 2007) onboard *Suzaku* (Mitsuda et al. 2007), in order to make clear the origin of the excess. The XIS is an ideal instrument to study the emission lines from the WHIM, because it does not show a large low-energy tail in the pulse-height distribution function, even in the very soft ($E \lesssim 0.5 \text{ keV}$) band.

We assume a Hubble constant of $70 \text{ km s}^{-1} \text{ Mpc}^{-1}$ or $h_{70} = 1$ and $\Omega_m = 0.3$, $\Omega_\Lambda = 0.7$. The solar metal abundance is given by Lodders (2003). Un-

Electronic address: Y.Takei@sron.nl

¹ SRON Netherlands Institute for Space Research, Sorbonnelaan 2, 3584CA, Utrecht, The Netherlands

² Massachusetts Institute of Technology, Kavli Institute for Astrophysics and Space Science, 77 Massachusetts Ave 37-551, Cambridge, MA 02139

³ University of Michigan, Department of Astronomy, 500 Church St., Ann Arbor, MI 48109

⁴ Institute of Space and Astronautical Science (ISAS), Japan Aerospace Exploration Agency (JAXA), 3-1-1 Yoshinodai, Sagamihara, Kanagawa, 229-8510, Japan

⁵ Department of Physics, Tokyo Metropolitan University, 1-1 Minami-Osawa, Hachioji, Tokyo 192-0397, Japan

⁶ Department of Physics, Kanazawa University, Kakuma-machi, Kanazawa 920-1192, Japan

TABLE 1
OBSERVATION INFORMATION

	Coordinates (J2000.0)	Exposure ^a
Coma-11	(12 ^h 58 ^m 31.7 ^s , 28°24′13.2″)	53.02 ks
Coma-7	(12 ^h 57 ^m 22.9 ^s , 28°08′58.1″)	25.26 ks
ComaBKG	(13 ^h 15 ^m 00.0 ^s , 31°39′28.0″)	30.78 ks

^a After screening as described in the text

less otherwise stated, errors and upper limits are at the 68 % confidence level in the figures, and at the 90 % confidence level in the text and tables. The emission line intensities are quoted in line units (LU) defined as photons cm⁻² s⁻¹ sr⁻¹. One LU corresponds to 8.46×10^{-8} photons cm⁻² s⁻¹ arcmin⁻².

2. OBSERVATIONS AND DATA REDUCTION

We observed three fields from 2007 Jun 19 to 2007 Jun 22; two of them are in the Coma outskirts, namely Coma-11 and Coma-7 that FBH03 observed with *XMM-Newton*, and the other is an offset observation (hereafter we call it ComaBKG) of 5.0-degree distance from the Coma center (NGC 4874). The coordinates and net exposure times are summarized in Table 1. The XIS was operated in the normal mode and with the spaced-row charge injection (SCI). We analysed the clean-event data of Ver 2.0 processing with standard parameters, but with the latest gain calibration (Ver 2.6.1.16). We excised the last 6.7 ks of the Coma-7 observation, during which the XIS suffered a problem with zero-level determination.

We excluded the regions where point sources are found or which are irradiated by the ⁵⁵Fe calibration sources. The point sources were identified using the *XMM-Newton* Serendipitous Source Catalogue (2XMM; Watson et al. 2008), as well as by examining the XIS images by eyes. Response files were created using *Suzaku* ftools xisrmfgen version 2007-05-14 and xissimarfgen version 2007-09-22 with CALDB released on 2008-01-14. We assumed uniform emission from the sky. The spectra and response files of the two XIS-FIs (XIS0 and 3) are added using ftools mathpha, marfmr, and addrmf. X-ray spectra are analysed with XSPEC 12.4.0.

3. SPECTRAL ANALYSIS

3.1. ComaBKG field

ComaBKG is observed in order to precisely determine the foreground and background emission, in particular O and Ne line intensities around the Coma cluster. The instrumental background, estimated using a ftool xisnxbgen version 2007-11-23 with COR2 as the sorting parameter (Tawa et al. 2008), were subtracted from the ComaBKG spectra.

We modeled the spectra with a sum of two thin thermal plasma models (APEC in XSPEC) and a power-law to represent emission from the local hot bubble (LHB), Milky-Way halo (MWH) and extragalactic cosmic X-ray background (CXB), respectively. MWH and CXB were convolved with absorption in the Galaxy (WABS in XSPEC). We fitted the spectra in the energy range 0.5–5.0 keV for the XIS-FI and 0.35–5.0 keV for XIS1 (XIS-BI). The spectra of XIS-FI and XIS-BI were fitted simultaneously. We ignored 1.40–1.55 keV where a

TABLE 2
BEST-FIT PARAMETERS OF COMABKG.

Parameter	Unit	Value
Galactic N_H	cm ⁻²	1.0×10^{20} (fixed)
LHB temperature	keV	0.07 (fixed)
LHB emission measure	pc cm ⁻⁶	$(3.0 \pm 3.0) \times 10^{-2}$
MWH temperature	keV	$0.144^{+0.021}_{-0.027}$
MWH emission measure	pc cm ⁻⁶	$(8.5^{+7.8}_{-3.0}) \times 10^{-3}$
CXB photon index		1.52 ± 0.09
CXB normalization ^a		7.7 ± 0.5
χ^2/dof thermal model		153.40/113=1.36
O VII intensity	LU	6.5 ± 1.0
O VIII intensity	LU	$1.1^{+0.9}_{-0.5}$
Ne IX intensity	LU	$0.15^{+0.15}_{-0.14}$
Ne X intensity	LU	$0 (< 0.2)$
χ^2/dof with lines		140.22/105=1.34

^a In a unit of photons cm⁻² s⁻¹ keV⁻¹ sr⁻¹ at 1 keV




FIG. 1.— Spectra and best-fit models (top) and residuals in units of sigma (bottom) of ComaBKG. Red and black show BI and FI, respectively. Instrumental background is subtracted.

relatively large uncertainty exists in the instrument background (Tawa et al. 2008). The redshift and abundance of the two thermal plasma models were fixed to 0 and 1.0 solar, respectively. The temperature of the LHB was also fixed to 0.07 keV, while the temperature of the MWH was allowed to vary. The column density of the Galactic hydrogen gas was fixed to 1.0×10^{20} cm⁻² (Kalberla et al. 2005). Figure 1 and Table 2 show the best-fit model.

We also fitted the spectra with another model in order to determine O VII, O VIII, Ne IX and Ne X intensity more precisely. Four Gaussians were added to represent the four line emissions, while O and Ne abundance of the LHB and MWH were fixed to zero. The energies of the Gaussians were allowed to vary ± 35 eV around the resonance line energy (0.574 keV for O VII, 0.653 keV for O VIII, 0.922 keV for Ne IX and 1.022 keV for Ne X) in order to compensate the uncertainty in the energy-scale calibration, while we fixed the line width to zero. The intensities of the four lines are also shown in Table 2. The relatively large χ^2 value may be due to incorrect calibration of the energy resolution. The χ^2 improved to 128.18 when we added 50 eV (FWHM) additional width into the model with lines. The best-fit parameters were not sensitive to this calibration uncertainty; they stay the same within the error at 90% confidence level. We confirmed that the fitting is not improved when we allow N_H to vary.

TABLE 3
BEST-FIT PARAMETERS OF COMA11 AND COMA7.

Parameter	Unit	Coma-11	Coma-7
Galactic N_{H}	cm^{-2}	1.0×10^{20} (fixed)	1.0×10^{20} (fixed)
Temperature	keV	$6.7^{+0.4}_{-0.3}$	$8.3^{+0.8}_{-0.6}$
Abundance	solar	0.36 ± 0.11	0.21 ± 0.15
Redshift		0.0231 (fixed)	0.0231 (fixed)
EM	pc cm^{-6}	0.100 ± 0.002	0.110 ± 0.002
χ^2/dof without lines		307.51/275=1.12	295.40/281=1.05
O VII intensity	LU	0.0 (< 0.9)	0.2 (< 2.1)
O VIII intensity	LU	1.1 ± 0.8	1.7 ± 1.1
Ne IX intensity	LU	0.1 (< 0.4)	0.4 (< 0.8)
Ne X intensity	LU	0.3 ± 0.2	0.3 (< 0.6)
χ^2/dof with lines		297.71/271=1.10	284.92/277=1.03




FIG. 2.— Spectra of Coma-11 (left) and Coma-7 (right). Red and black data correspond to BI and FI respectively. ComaBKG spectra were subtracted to correct the background emission.

3.2. Coma-11 and Coma-7 fields

The back- and foreground emission of Coma-11 and Coma-7 were corrected by subtracting the ComaBKG spectra that were extracted from the same region as Coma-11 or Coma-7 in the detector coordinates. We then fitted the spectra 0.5–7.5 keV for XIS-FI and 0.35–7.5 keV for XIS-BI by the emission of the intracluster medium (ICM), i.e., a thin thermal plasma model (APEC) convolved with Galactic absorption (wabs). The spectra of XIS-FI and XIS-BI were again fitted together. The spectra and the best-fit models are shown in Figure 2, and the best-fit parameters are in Table 3. We did not see a soft excess, i.e., emission from the WHIM. Next, we added four Gaussians to the model, corresponding to lines of O VII, O VIII, Ne IX and Ne X at the cluster redshift, in order to investigate possible contributions of the WHIM. The central energy of each line was fixed to that of the resonance lines at redshift $z = 0.0231$, and the width of the line was fixed to zero. The results are shown in Table 3. The fit is slightly improved ($\Delta\chi^2 \sim 10$) by adding the lines. However, the significance of each line is less than 3σ . The fitting without FI also gave the same results within the statistical errors, which excludes the possible systematic bias due to the low statistics of the FI data below 1 keV. We also tried to fit the spectra with two thin thermal plasma models, i.e., one for the intracluster medium and one for the WHIM. This did not improve the fit. To obtain robust upper limits of line intensities, we consider systematic uncertainties in calibration: the centroid energy (± 35 eV), the energy res-

olution (10–80 eV FWHM), energy scale (± 50 eV), and the contamination thickness ($\pm 5 \times 10^{-17} \text{ cm}^{-2}$). The 2σ upper limits we obtained for O VII, O VIII, Ne IX, and Ne X are 1.4, 2.7, 0.5, and 0.9 LU for Coma-11 and 2.6, 3.7, 1.2, and 1.1 LU for Coma-7, respectively. The uncertainty in Galactic O intensity should also be taken into account. From the variation among background fields in T07, we adopted ± 1.2 LU for O VII and ± 0.6 LU for O VIII as 1σ . Then, O VII and O VIII 2σ upper limits are 2.8 and 2.9 LU for Coma-11 and 3.5 and 3.9 LU for Coma-7, respectively.

4. DISCUSSION

We measured the spectra of Coma-11 and Coma-7 with the Suzaku XIS. The spectra were fitted with a single thin thermal plasma model and the emission from the WHIM was not confirmed. Here we compare our results with previous *XMM-Newton* observations. We do not discuss the lower energy excess ($E < 0.5$ keV) observed with ROSAT and EUVE, because the XIS is not sensitive in the energy.

FBH03 observed Coma-11 and Coma-7 with *XMM-Newton*. The O VII and O VIII intensity measured in FBH03 are 6.4 ± 0.8 and 6.2 ± 0.8 LU in Coma-11, and 1.4 ± 0.8 and 0.9 ± 0.6 LU in Coma-7, respectively. The intensities of Coma-7 in FBH03 are consistent with our *Suzaku* observation, while those of Coma-11 are discrepant at the 5σ level. The best-fit ICM temperature and abundance in Coma-11 are also inconsistent. Even if we fixed the temperature and abundance to the best-fit




FIG. 3.— Solar wind proton flux as a function of time, around Coma-11 and Coma-7 observations with *XMM-Newton* (left) and around our observations with *Suzaku* (right), measured with ACE SWEPAM instrument. The time of *XMM-Newton* and *Suzaku* observations is indicated with vertical lines.

values of FBH03, the discrepancies of the O line intensities remain as large. The O intensities observed with *Suzaku* are similar to those of Coma-7; we did not confirm the spatial variation.

T07 measured O VII, O VIII and Ne IX emission line intensities with *XMM-Newton* in the field of X Com and four background fields whose distance from NGC 4874 is 46.2–162.7 arcmin. The intensities of O VII, O VIII and Ne IX lines measured by T07 are 6.6 ± 0.2 , 1.8 ± 0.2 , and 0.7 ± 0.1 LU in the X Com field and 7.4 ± 1.2 , 2.6 ± 0.6 , and 0.4 ± 0.1 LU, in the (averaged) background fields, respectively. The values of the X Com field shown above contain contributions of the background emission. The intensities of the X Com field are consistent within 90% confidence range with those of the Coma-11 and Coma-7 measured in this work, and those of the background field of T07 are also consistent with those of ComaBKG.

The origin of the large discrepancy in Coma-11 O intensities between this work and FBH03 could be contamination by SWCX emission. The solar wind proton flux observed with ACE (Advanced Composition Explorer⁷) showed flares both during Coma-11 observation of FBH03 ($2.5 \times 10^9 \text{ cm}^{-2} \text{ s}^{-1}$) and during ComaBKG observation of this work ($1.5 \times 10^9 \text{ cm}^{-2} \text{ s}^{-1}$) as shown in Figure 3. Both are ~ 8 times larger than the quiescent state around each observation time. Hence, two possibilities should be considered: O intensities of Coma-11 in FBH03 are overestimated, or those of ComaBKG in this work are overestimated and thus those of Coma-11 in this work are underestimated. However, the latter case is not likely because the O-line intensities are consistent with or even smaller than T07 measurements of background fields around the Coma cluster. There is no indication of the geocoronal SWCX emission; the first point in the line of sight (LOS) where the geomagnetic field started open was always higher than 10 times Earth radius⁸ (Fujimoto et al. 2007; Tsyganenko & Sitnov 2005), and there was no variation in the O VII and O VIII flux during the ComaBKG observation. The heliospheric SWCX emission could be weak since it is faint at solar minimum on the high ecliptic latitude line of sight (Koutroumpa et al. 2007). Moreover, even in the case that the ComaBKG observation was heavily contami-

nated with the SWCX emission, the 2σ upper limits of Coma-11 O VII and O VIII intensity would be no larger than 4.8 LU and 3.5 LU, respectively, which are still lower than values of FBH03. These values are calculated by assuming the lower limits of O VII and O VIII intensities as the LHB contribution without any MWH contribution, measured toward a molecular cloud MBM 12 (3.5 LU and 0 LU, respectively; Smith et al. 2007). On the other hand, it is more likely that the FBH03 observation suffered heliospheric SWCX contamination, since a similar solar wind flare during *XMM-Newton* observation of the Hubble deep field north increased the O VII and O VIII flux by 6.25 ± 0.67 and 5.53 ± 2.87 LU, respectively (Snowden et al. 2004; Koutroumpa et al. 2007). The solar cycle at the observations, the position of the Earth and the LOS, which also affects the X-ray flux induced by SWCX, are similar between the two observations. Note that the observation of FBH03 was performed at solar maximum, while our observation was at solar minimum. The field-to-field variation of O VII and O VIII flux reported by FBH03 might also be due to the variation of the SWCX flux.

YT and SK thank T. Hagihara for his instruction on the calculation of the geomagnetic field distribution. YT is financially supported by the Dutch Organization for Scientific Research (NWO). EDM and JNB gratefully acknowledge support for *Suzaku* activities from NASA.

⁷ <http://www.srl.caltech.edu/ACE/ASC>

⁸ Calculated using the software GEOPACK-2005 and T96 magnetic field model (<http://geo.phys.spbu.ru/~tsyganenko/modeling.html>), with CDAWeb solar-wind parameters (http://cdaweb.gsfc.nasa.gov/cdaweb/sp_phys/)

REFERENCES

- Bonamente, M., Joy, M. K., & Lieu, R. 2003, *ApJ*, 585, 722
- Bonamente, M., Lieu, R., & Kaastra, J. 2005, *A&A*, 443, 29
- Bregman, J. N. 2007, *ARA&A*, 45, 221
- Bregman, J. N., & Lloyd-Davies, E. J. 2006, *ApJ*, 644, 167
- Cen, R., & Ostriker, J. P. 2006, *ApJ*, 650, 560
- Cravens, T. E. 2000, *ApJ*, 532, L153
- Danforth, C. W., & Shull, J. M. 2007, *astro-ph/0709.4030*
- Finoguenov, A., Briel, U. G., & Henry, J. P. 2003, *A&A*, 410, 777
- Fujimoto, R., Mitsuda, K., McCammon, D., Takei, Y., Bauer, M., Ishisaki, Y., Porter, S. F., Yamaguchi, H., Hayashida, K., & Yamasaki, N. Y. 2007, *PASJ*, 59, 133
- Fukugita, M., Hogan, C. J., & Peebles, P. J. E. 1998, *ApJ*, 503, 518
- Kaastra, J. S., Lieu, R., Tamura, T., Paerels, F. B. S., & den Herder, J. W. 2003, *A&A*, 397, 445
- Kaastra, J. S., Werner, N., Herder, J. W. A. d., Paerels, F. B. S., de Plaa, J., Rasmussen, A. P., & de Vries, C. P. 2006, *ApJ*, 652, 189
- Kalberla, P. M. W., Burton, W. B., Hartmann, D., Arnal, E. M., Bajaja, E., Morras, R., & Pöppel, W. G. L. 2005, *A&A*, 440, 775
- Koutroumpa, D., Acero, F., Lallement, R., Ballet, J., & Kharchenko, V. 2007, *A&A*, 475, 901
- Koyama, K., Tsunemi, H., Dotani, T., Bautz, M. W., Hayashida, K., Tsuru, T. G., Matsumoto, H., Ogawara, Y., Ricker, G. R., Doty, J., Kissel, S. E., Foster, R., Nakajima, H., Yamaguchi, H., Mori, H., Sakano, M., Hamaguchi, K., Nishiuchi, M., Miyata, E., Torii, K., Namiki, M., Katsuda, S., Matsuura, D., Miyauchi, T., Anabuki, N., Tawa, N., Ozaki, M., Murakami, H., Maeda, Y., Ichikawa, Y., Prigozhin, G. Y., Boughan, E. A., Lamarr, B., Miller, E. D., Burke, B. E., Gregory, J. A., Pillsbury, A., Bamba, A., Hiraga, J. S., Senda, A., Katayama, H., Kitamoto, S., Tsujimoto, M., Kohmura, T., Tsuboi, Y., & Awaki, H. 2007, *PASJ*, 59, 23
- Lodders, K. 2003, *ApJ*, 591, 1220
- Mitsuda, K., Bautz, M., Inoue, H., Kelley, R. L., Koyama, K., Kunieda, H., Makishima, K., Ogawara, Y., Petre, R., Takahashi, T., Tsunemi, H., White, N. E., Anabuki, N., Angelini, L., Arnaud, K., Awaki, H., Bamba, A., Boyce, K., Brown, G. V., Chan, K.-W., Cottam, J., Dotani, T., Doty, J., Ebisawa, K., Ezoe, Y., Fabian, A. C., Figueroa, E., Fujimoto, R., Fukazawa, Y., Furusho, T., Furuzawa, A., Gendreau, K., Griffiths, R. E., Haba, Y., Hamaguchi, K., Harrus, I., Hasinger, G., Hatsukade, I., Hayashida, K., Henry, P. J., Hiraga, J. S., Holt, S. S., Hornschemeier, A., Hughes, J. P., Hwang, U., Ishida, M., Ishisaki, Y., Isobe, N., Itoh, M., Iyomoto, N., Kahn, S. M., Kamae, T., Katagiri, H., Kataoka, J., Katayama, H., Kawai, N., Kilbourne, C., Kinugasa, K., Kissel, S., Kitamoto, S., Kohama, M., Kohmura, T., Kokubun, M., Kotani, T., Kotoku, J., Kubota, A., Madejski, G. M., Maeda, Y., Makino, F., Markowitz, A., Matsumoto, C., Matsumoto, H., Matsuoka, M., Matsushita, K., McCammon, D., Mihara, T., Misaki, K., Miyata, E., Mizuno, T., Mori, K., Mori, H., Morii, M., Moseley, H., Mukai, K., Murakami, H., Murakami, T., Mushotzky, R., Nagase, F., Namiki, M., Negoro, H., Nakazawa, K., Nousek, J. A., Okajima, T., Ogasaka, Y., Ohashi, T., Oshima, T., Ota, N., Ozaki, M., Ozawa, H., Parmar, A. N., Pence, W. D., Porter, F. S., Reeves, J. N., Ricker, G. R., Sakurai, I., Sanders, W. T., Senda, A., Serlemitsos, P., Shibata, R., Soong, Y., Smith, R., Suzuki, M., Szymkowiak, A. E., Takahashi, H., Tamagawa, T., Tamura, K., Tamura, T., Tanaka, Y., Tashiro, M., Tawara, Y., Terada, Y., Terashima, Y., Tomida, H., Torii, K., Tsuboi, Y., Tsujimoto, M., Tsuru, T. G., Turner, M. J. L., Ueda, Y., Ueno, S., Ueno, M., Uno, S., Urata, Y., Watanabe, S., Yamamoto, N., Yamaoka, K., Yamasaki, N. Y., Yamashita, K., Yamauchi, M., Yamauchi, S., Yaqoob, T., Yonetoku, D., & Yoshida, A. 2007, *PASJ*, 59, 1
- Nicastro, F., Mathur, S., Elvis, M., Drake, J., Fang, T., Fruscione, A., Krongold, Y., Marshall, H., Williams, R., & Zezas, A. 2005, *Nature*, 433, 495
- Rasmussen, A. P., Kahn, S. M., Paerels, F., Herder, J. W. d., Kaastra, J., & de Vries, C. 2007, *ApJ*, 656, 129
- Rauch, M., Miralda-Escude, J., Sargent, W. L. W., Barlow, T. A., Weinberg, D. H., Hernquist, L., Katz, N., Cen, R., & Ostriker, J. P. 1997, *ApJ*, 489, 7
- Smith, R. K., Bautz, M. W., Edgar, R. J., Fujimoto, R., Hamaguchi, K., Hughes, J. P., Ishida, M., Kelley, R., Kilbourne, C. A., Kuntz, K. D., McCammon, D., Miller, E., Mitsuda, K., Mukai, K., Plucinsky, P. P., Porter, F. S., Snowden, S. L., Takei, Y., Terada, Y., Tsuboi, Y., & Yamasaki, N. Y. 2007, *PASJ*, 59, 141
- Snowden, S. L., Collier, M. R., & Kuntz, K. D. 2004, *ApJ*, 610, 1182
- Spiegel, D. N., Bean, R., Doré, O., Nolte, M. R., Bennett, C. L., Dunkley, J., Hinshaw, G., Jarosik, N., Komatsu, E., Page, L., Peiris, H. V., Verde, L., Halpern, M., Hill, R. S., Kogut, A., Limon, M., Meyer, S. S., Odegard, N., Tucker, G. S., Weiland, J. L., Wollack, E., & Wright, E. L. 2007, *ApJS*, 170, 377
- Takei, Y., Henry, J. P., Finoguenov, A., Mitsuda, K., Tamura, T., Fujimoto, R., & Briel, U. G. 2007, *ApJ*, 655, 831
- Tawa, N., Hayashida, K., Nagai, M., Nakamoto, H., Tsunemi, H., Yamaguchi, H., Ishisaki, Y., Miller, E. D., Mizuno, T., Dotani, T., Ozaki, M., & Katayama, H. 2008, *PASJ*, 60, 11
- Tsyganenko, N. A., & Sitnov, M. I. 2005, *Journal of Geophysical Research (Space Physics)*, 110, 3208
- Watson, M., & et al. 2008 in prep.